

Appendix H. Forestry & Land Use

Overview

Forestland emissions refer to the net carbon dioxide (CO₂) flux¹ from forested lands in Florida, which account for about 47% of the state's land area.² The dominant forest type in Florida is Longleaf-slash pine which makes up about 35.6% of forested lands. Other common forest types are Oak-gum-cypress at 26.18%, Oak-hickory at 17.3%, ~~Loblolly-shortleaf at 9%~~, and ~~at~~ Oak-pine at 9% of forested land. All other forest types make up less than 4.6% each of the State's forests.

Through photosynthesis, CO₂ is taken up by trees and plants and converted to carbon in biomass within the forests. Carbon dioxide emissions occur from respiration in live trees, decay of dead biomass, and combustion (both wildland fires and biomass removed from forests for energy use). In addition, carbon is stored for long time periods when forest biomass is harvested for use in durable wood products. Carbon dioxide flux is the net balance of CO₂ removals from and emissions to the atmosphere from the processes described above.

The forestry sector ~~net GHG emissions are~~ ~~CO₂ flux is~~ categorized into two primary subsectors:

- *Forested Landscape*: this consists of carbon flux occurring on lands that are not part of the urban landscape. Fluxes covered include net carbon sequestration, carbon stored in harvested wood products (HWP) or landfills, and emissions from forest fires.
- *Urban Forestry and Land Use*: this covers carbon sequestration in urban trees, carbon flux associated with carbon storage from landscape waste and food scraps in landfills, and nitrous oxide (N₂O) emissions from settlement soils (those occurring as a result of application of synthetic fertilizers in urbanized areas).

Inventory and Reference Case Projections

Forested Landscape

For over a decade, the United States Forest Service (USFS) has been developing and refining a forest carbon modeling system for the purposes of estimating forest carbon inventories. The methodology is used to develop national forest CO₂ fluxes for the official *US Inventory of Greenhouse Gas Emissions and Sinks*. The national estimates are compiled from state-level data. The Florida forest CO₂ flux data in this report come from the national analysis and are provided by the USFS. See the footnotes below for the most current documentation for the forest carbon modeling.³ Additional forest carbon information is in the form of specific carbon conversion factors.⁴

¹ "Flux" refers to both emissions of CO₂ to the atmosphere and removal (sinks) of CO₂ from the atmosphere.

² Total forested acreage is 16.12 million acres in 2005. Acreage by forest type available from the USFS Southern Research Station report, *Florida's Forests – 2005 Update* at: <http://www.treearch.fs.fed.us/pubs/28996> <http://www.fs.fed.us/ne/global/pubs/books/epa/states/FL.htm>. The total land area in Florida is 34.6 million acres (<http://www.50states.com/Florida.htm>).

³ The most current citation for an overview of how the USFS calculates the inventory based forest carbon estimates as well as carbon in harvested wood products is from the US Inventory of Greenhouse Gas Emissions and Sinks: 1990-2005 (and earlier editions), US Environmental Protection Agency, Report # USEPA #430-R-07-002, April 2007, available at: <http://epa.gov/climatechange/emissions/usinventoryreport.html>. Both Annex 3.12 and Chapter 7

The forest CO₂ flux methodology relies on input data in the form of plot-level forest volume statistics from the Forest Inventory and Analysis (FIA) Program. FIA data on forest volumes are converted to values for ecosystem carbon stocks (i.e., the amount of carbon stored in forest carbon pools) using the FORCARB2 modeling system. Coefficients from FORCARB2 are applied to the plot level survey data to give estimates of C density [megagrams (Mg) per hectare] for a number of separate C pools (see Table H1 for Florida C pools). Additional background on the FORCARB system is provided in a number of publications.⁵

Carbon dioxide flux is estimated as the change in carbon mass for each carbon pool over a specified time-frame. Forest biomass data from at least two points in time are required. The change in carbon stocks between time intervals is estimated for specific carbon pools (Live Tree, Standing Dead Wood, Understory, Down & Dead Wood, Forest Floor, and Soil Organic Carbon) and divided by the number of years between inventory samples. Annual increases in carbon density reflect carbon sequestration in a specific pool; decreases in carbon density reveal CO₂ emissions or carbon transfers out of that pool (e.g., death of a standing tree transfers carbon from the live tree to standing dead wood pool). The amount of carbon in each pool is also influenced by changes in forest area (e.g., an increase in area could lead to an increase in the associated forest carbon pools and the estimated flux). The sum of carbon stock changes for all forest carbon pools yields a total net CO₂ flux for forest ecosystems.

In preparing these estimates, USFS estimates the amount of forest carbon in different forest types as well as different carbon pools. The different forests also include differences in ownership class: those in the national forest (NF) system and those that are not federally-owned (private and other public forests). Additional details on the forest carbon inventory methods can be found in Annex 3 to the US EPA's 2007 GHG inventory for the US.⁶

Carbon pool data for three FIA cycles to estimate flux for two different periods were available for Florida. The carbon pool data for three points in time are shown in Table H1 below. Note that prior to 1995, the Southern FIA Program took periodic forest inventory surveys for Florida (approximately on a 10-year schedule). Beginning in 2001, Florida transitioned from periodic to annual inventories as modifications to the FIA program were applied. The annual inventory measures is supposed to measure 20 percent% of the plots in Florida each year, and delivers a

LULUCF are useful sources of reference. See also Smith, J.E., L.S. Heath, and M.C. Nichols (in press), *US Forest Carbon Calculation Tool User's Guide: Forestland Carbon Stocks and Net Annual Stock Change*, Gen Tech Report, Newtown Square, PA: US Department of Agriculture, Forest Service, Northern Research Station.

⁴ Smith, J.E., and L.S. Heath (2002). "A model of forest floor carbon mass for United States forest types," Res. Pap. NE-722. Newtown Square, PA: US Department of Agriculture, Forest Service, Northeastern Research Station. 37 p., or Jenkins, J.C., D.C. Chojnacky, L.S. Heath, R.A. Birdsey (2003), "National-scale biomass estimators for United States tree species", *Forest Science*, 49:12-35.

⁵ Smith, J.E., L.S. Heath, and P.B. Woodbury (2004). "How to estimate forest carbon for large areas from inventory data", *Journal of Forestry*, 102: 25-31; Heath, L.S., J.E. Smith, and R.A. Birdsey (2003), "Carbon trends in US forest lands: A context for the role of soils in forest carbon sequestration", In J. M. Kimble, L. S. Heath, R. A. Birdsey, and R. Lal, editors. *The Potential of US Forest Soils to Sequester Carbon and Mitigate the Greenhouse Effect*. CRC Press, New York; and Woodbury, Peter B.; Smith, James E.; Heath, Linda S. 2007, "Carbon sequestration in the US forest sector from 1990 to 2010", *Forest Ecology and Management*, 241:14-27.

⁶ Annex 3 to EPA's 2007 report, which contains estimates for calendar year 2005, can be downloaded at: <http://www.epa.gov/climatechange/emissions/downloads06/07Annex3.pdf>.

complete inventory report every 5 years. The 2005 inventory, however, only covers 60 percent of all sample plots in the State, due to a slower start of the annual inventory program, and delivers a complete inventory report every 5 years. Florida completed its first annual inventory cycle in 2005. According to USFS, 20 percent of the sample plots were scheduled to be completed annually after 2005.

These underlying FIA data, as shown in Table H1, display a net decrease of 402,000 acres in forested area between 1987 and 2005. Between 1987 and 1995, forested area decreased a significant 663,328,000 acres; forested area increased an additional 264,74,000 acres from 1995 to 2005. Most of the forested lands in Florida are considered timberland, meaning they are unreserved productive forestland producing, or capable of producing, crops of industrial wood. The timberland area is shown to have decreased by 332,000 acres between 1987 and 1995 while it increased 901,000 acres between 1995 and 2005. This increase in timberland area kept total carbon levels in 2005 (1,100 million metric tons) similar to the carbon level in 1987 (1,116 million metric tons) despite overall forest area decreases. It is not clear based on currently available information how much of this growth in forested-timberland area is due to methodological changes (see Key Uncertainties section below) or land use conversion (e.g. agricultural use to forested use).

Table H1. USFS Forest Carbon Pool Data for Florida

Forest Pool	1987 (MMtC)	1995 (MMtC)	2005 (MMtC)
Live Tree—Above Ground	242	236	275
Live Tree—Below Ground	48.9	47.8	55.4
Understory	21.1	20.6	20.5
Standing Dead	11.4	11.2	11.1
Down Dead	19.0	18.7	22.2
Forest Floor	51.4	49.5	51.5
Soil Carbon	722	662	664
Totals	1,116	1,046	1,100
Forest Area	1987 (10 ³ -acres)	1995 (10 ³ -acres)	2005 (10 ³ -acres)
All Forests	16,549	15,866	16,147
Timberland	14,983	14,651	15,552

Forest Pool	1987 (MMtC)	Adjusted 1995 (MMtC)	2005 (MMtC)
Live Tree – Above Ground	241.9	241.5	274.9

Formatted: Font: 10 pt

Formatted Table

Formatted: Font: 10 pt

<u>Live Tree – Below Ground</u>	48.9	48.9	55.4
<u>Understory</u>	21.1	21.1	20.5
<u>Standing Dead</u>	11.4	11.4	11.1
<u>Down Dead</u>	19.0	19.2	22.2
<u>Forest Floor</u>	51.4	50.6	51.5
<u>Soil Carbon</u>	722.4	676.7	664.2
Totals	1,116	1,069	1,100
Forest Area	1987 (10³ acres)	1995 (10³ acres)	2005 (10³ acres)
<u>All Forests</u>	16,549	16,221	16,147
<u>Timberland</u>	14,983	14,651	15,552

MMtC = million metric tons of carbon. Positive numbers indicate net emission. Multiply MMtC by 3.67 (44/12) to convert to MMtCO₂.

Totals may not sum exactly due to independent rounding.

Data source: Smith, James, et al. *US Forest Carbon Calculation Tool: Forest-Land Carbon Stocks and Net Annual Stock Change* (<http://www.nrs.fs.fed.us/pubs/2394>), December 2007. Adjusted for 1995 total Forest Area: Brown, Mark, J. *Florida's Forests – 2005 Update* (<http://www.treesearch.fs.fed.us/pubs/28996>), June 2008.

Table H2 shows the annualized carbon stocks interpolated from Florida FIA data using the Carbon Calculation Tool (CCT).⁷ These annualized carbon stocks differ from the carbon stocks in Table H1 in that they are interpolated values (between forest inventory years) to January 1st of each year. The difference in carbon between each consecutive year is the carbon flux for that year. The carbon fluxes for each period shown in Table H3 are based on these annualized carbon stock estimates.

Table H2. Annualized Forest Carbon Pools from USFS Carbon Calculation Tool

Forest Pool	1990 (MMtC)	1995 (MMtC)	2005 (MMtC)
Live Tree – Above Ground	240	238	281
Live Tree – Below Ground	48.5	48.3	56.5
Understory	20.9	20.6	20.5
Standing Dead	11.3	11.2	11.1
Down Dead	18.9	18.9	22.7
Forest Floor	50.7	49.6	51.8
Soil Carbon	699	662	665
Totals	1,090	1,049	1,108
Forest Area	1990 (10³ acres)	1995 (10³ acres)	2005 (10³ acres)

⁷ Smith, James, et al. *US Forest Carbon Calculation Tool: Forest-Land Carbon Stocks and Net Annual Stock Change* (<http://www.nrs.fs.fed.us/pubs/2394>), November 2007.

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: Italic

Formatted: Font: 10 pt, No underline, Font color: Auto

Formatted: Font: 10 pt, No underline, Font color: Auto

All Forests	16,290	15,881	16,188
Timberland	14,857	14,700	15,684

In addition to the forest carbon pools, additional carbon is stored in biomass removed from the forest for the production of harvested wood products (HWP). Carbon remains stored in the durable wood products pool or is transferred to landfills where much of the carbon remains stored over a long period of time. The USFS uses a model referred to as WOODCARB2 for the purposes of modeling national HWP carbon storage.⁸ State-level information for Florida was provided to CCS by USFS.⁹

As shown in Table H2, about 3.9 million metric tons (MMt) of CO₂ per year (yr) is estimated by the USFS to be sequestered annually (1990-2005) in wood products. Also, as shown in this table, the total flux estimate including all forest pools is ~~31.5~~19.7 MMtCO₂e/yr between 1987 and 1995, and ~~-25.4~~-16.0 MMtCO₂e/yr between 1995 and 2005.¹⁰ This fluctuation is largely due to significant differences in forest carbon pools from each cycle period (note the differences in Table H1₂ on forested area between these two periods), as well as the fluctuation in the soil organic carbon pool. ~~Note that from 1995 to 2005, soil carbon was considered a net sink, in addition to each all of the forest carbon pools collectively.~~ Given the changes noted above in timberland, it appears that much of the negative trend in carbon flux (sequestration) is from the increase in timberland between 1995 and 2005.

Table H2. USFS Annual Forest Carbon Fluxes for Florida

<u>FL Forest Pool</u>	<u>1987-1995 Flux (MMtCO₂)</u>	<u>1995-2005 Flux (MMtCO₂)</u>
<u>Live Tree</u>	0.19	-16.0
<u>Understory</u>	0.02	0.23
<u>Standing Dead</u>	-0.03	0.14
<u>Down Dead</u>	-0.08	-1.21
<u>Forest Floor</u>	0.40	-0.35
<u>Soil Carbon</u>	23.1	5.01
<u>Harvested Wood Products</u>	-3.89	-3.89
<u>Totals</u>	19.7	-16.0
<u>Totals (excluding soil carbon)</u>	-3.38	-21.1

Formatted Table

⁸ Skog, K.E., and G.A. Nicholson (1998), "Carbon cycling through wood products: the role of wood and paper products in carbon sequestration", *Forest Products Journal*, 48, (7/8):75-83; or Skog, K.E., K. Pingoud, and J.E. Smith (2004), "A method countries can use to estimate changes in carbon stored in harvested wood products and the uncertainty of such estimates", *Environmental Management*, 33, (Suppl. 1): S65-S73.

⁹ Obtained from the Harvested Wood Product model developed by Ken Skog, USFS

¹⁰ Jim Smith, USFS, *US Forest Carbon Calculation Tool: Forest-Land Carbon Stocks and Net Annual Stock Change* (<http://www.nrs.fs.fed.us/pubs/2394>), December 2007.

Forest Pool	1987-1995 Flux (MMtCO ₂)	1995-2005 Flux (MMtCO ₂)
Forest Carbon Pools (non-soil)	4.85	-20.6
Soil Organic Carbon	30.6	-0.92
Harvested Wood Products	-3.89	-3.89
Totals	31.5	-25.4
Totals (excluding soil carbon)	0.96	-24.5

Totals may not sum exactly due to independent rounding.
 Data source: Smith, James, et al. US Forest Carbon Calculation Tool: Forest-Land Carbon Stocks and Net Annual Stock Change (<http://www.nrs.fs.fed.us/pubs/2394>), USFS, December 2007.

Based on discussions with the USFS, CCS recommends excluding the soil carbon pool from the overall forest flux estimates due to a high level of uncertainty associated with these estimates. The forest carbon flux estimates provided in the summary tables at the front of this report are those without the soil carbon pool.

For historic emission estimates, CCS used the 1987-1995 carbon flux to represent yearly forest carbon flux prior to 1995. Current flux estimates (1995-2005) are from the 1995 inventory and 2005 annual inventory stocks. For the reference case projections (2005-2025), the total forest area is assumed to decrease at a rate of 7420 acres per year (the same rate of decrease as 1995-2005 total forest area). ~~and~~ The carbon densities of forestlands ~~are~~ were assumed to remain at the same levels as in 2005 since information is not available on the near term effects of climate change and their impacts on forest productivity. Table H3 shows the 2005-2025 reference case projections and the assumptions used for estimating the projected CO₂ flux. Nor were data readily available on projected losses/gains in forested area.

Table H3. Reference Case Projections (2005-2025) and Assumptions

	2005	2025
Total Forest Land (thousand acres)	16,147	15,999
Carbon Pool (MMtC)	1,100	1,090
Carbon Density (Mg/acre)	68.1	68.1
	1995-2005	2005-2025
Carbon Flux (MMtC/yr)	-5.7	-5.2
CO ₂ Flux (MMtCO ₂ /yr)	-21.1	-19.2

Formatted: Font: No underline, Font color: Auto, Subscript

Formatted: Caption, Keep with next

Formatted Table

Urban Forestry & Land Use

GHG emissions for 1990 through 2005 were estimated using the EPA State Inventory Tool (SIT) software and the methods provided in the Emission Inventory Improvement Program (EIIP) guidance document for the sector.¹¹ In general, the SIT methodology applies emission factors developed for the US to activity data for the urban forestry sector. Activity data include urban area, urban area with tree cover, amount of landfilled yard trimmings and food scraps, and the total amount of synthetic fertilizer applied to settlement soils (e.g., parks, yards, etc.). This methodology is based on international guidelines developed by sector experts for preparing GHG emissions inventories.¹² Table H43 displays the emissions and reference case projections for Florida.

Table H3H4. Urban Forestry Emissions and Reference Case Projections (MMtCO₂e)

Urban Forestry & Land Use Subsector	1990	2000	2005	2010	2020	2025
Urban Trees	-1.78	-2.31	-2.58	-2.58	-2.58	-2.58
Landfilled Yard Trimmings and Food Scraps	-12.8	-3.44	-3.75	-3.75	-3.75	-3.75
N ₂ O from Settlement Soils	0.1	0.1	0.1	0.1	0.1	0.1
Total	-14.4	-5.65	-6.23	-6.23	-6.23	-6.23

Formatted: Font: (Default) Times New Roman, No underline,
Formatted: Font: (Default) Times New Roman

*Data for settlement soils was obtained from AAPFCO (2006) Commercial Fertilizers 2005. Association of American Plant Food Control Officials and The Fertilizer Institute. University of Kentucky, Lexington, KY.

Changes in carbon stocks in urban trees are equivalent to tree growth minus biomass losses resulting from pruning and mortality. Net carbon sequestration was calculated using data on crown cover area. The default urban area data in SIT (which varied from 12,518 square kilometers [km²] to 18,131 km² between 1990 and 2005) was multiplied by the state estimate of the percent of urban area with tree cover (18% for Florida) to estimate the total area of urban tree cover. These default SIT urban area tree cover data represent area estimates taken from the US Census and coverage for years 1990 and 2000.¹³ Estimates of urban area in the intervening years (1990-1999) and subsequent years (2001-2005) are interpolated and extrapolated, respectively.

Estimates of net carbon flux of landfilled yard trimmings and food scraps were calculated by estimating the change in landfill carbon stocks between inventory years. Carbon stock estimates were calculated by determining the mass of landfilled carbon resulting from yard trimmings or food scraps discarded in a given year, adding the accumulated landfilled carbon from previous years, and subtracting the portion of carbon landfilled in previous years that decomposed.

¹¹ GHG emissions were calculated using SIT, with reference to EIIP, Volume VIII: Chapter 8.
¹² Revised 1996 Intergovernmental Panel on Climate Change (IPCC) Guidelines for National Greenhouse Gas Inventories, published by the National Greenhouse Gas Inventory Program of the IPCC, available at (<http://www.ipcc-nggip.iges.or.jp/public/gl/invs1.htm>); and Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories, published in 2000 by the National Greenhouse Gas Inventory Program of the IPCC, available at: (<http://www.ipcc-nggip.iges.or.jp/public/gp/english/>).
¹³ Dwyer, John F.; Nowak, David J.; Noble, Mary Heather; Sisinni, Susan M. 2000. Connecting people with ecosystems in the 21st century: an assessment of our nation's urban forests. Gen. Tech. Rep. PNW-GTR-490

Default SIT landfilled yard trimmings and food scraps data were estimated using the Florida State population and the national yard trimmings and food scraps ratio. Along with the national trend, Florida's landfilled yard trimmings and food scraps decreased significantly during the 1990's. This is largely due to programs discouraging or banning disposal and a dramatic increase in the number of municipal composting facilities, which reduced the proportion of collected yard trimmings that are discarded in landfills.¹⁴ This decrease in landfilled yard trimmings and food scraps disposal rate has resulted in a decrease in the rate of landfill carbon storage to 3.75 MMtCO₂e in 2005 from 12.8 MMtCO₂e in 1990.


Settlement soils include all developed land, transportation infrastructure and human settlements of any size. N₂O emissions from settlement soils were calculated in SIT using default synthetic fertilizer data multiplied by N₂O emission factor. Future projections of CO₂ fluxes from urban trees, landfilled yard trimmings and food scraps, and settlement soils were kept constant at 2005 levels. Table H4 provides a summary of the estimated flux for the entire forestry and land use sector.

Wildfire and Prescribed Burning Emissions

Biomass burned in forest fires emits CO₂, methane (CH₄), and N₂O, in addition to many other gases and pollutants. Since CO₂ emissions are captured under total carbon flux calculations in USFS' modeling described above, CCS used SIT to estimate CH₄ and N₂O emissions. No default data were available for area burned by forest type, so CCS used available state data from Florida Department of Agriculture & Consumer Services.¹⁵ Wildfire acres burned data were used for the years 1990-2007⁵ and the forest type of "other temperate forests" was assumed in SIT to calculate historical emissions. [Prescribed forest fires area burned data were available for 1998-2007 and were entered in SIT. Figure H1 compares the yearly wildfire and prescribed fire areas in Florida.](#)

¹⁴ INVENTORY OF U.S. GREENHOUSE GAS EMISSIONS AND SINKS: 1990-2005, Land Use, Land Use Change and Forestry (<http://www.epa.gov/climatechange/emissions/usinventoryreport.html>).

¹⁵ Wildfire acres burned data, and Authorized burning data obtained from Florida Department of Agriculture & Consumer Services, Division of Forestry, Wildland Fire (<http://tlhforweb1.doacs.state.fl.us/PublicReports/>), [December 2007/June 2008](#).



Formatted: Keep with next

Figure H1. Florida Fires Area Burned (Thousand Acres)

Formatted: Caption, Widow/Orphan control

Data were not available for prescribed fires between 1990 and 1997 so an average of 1998-2007 acres burned were assumed to estimate emissions for 1990-1997. Due to the yearly fluctuation of both wildfire and prescribed fire data, projected emissions for 2008-2025 were assumed to be the average of 1990-2007 wildfire emissions and 1998-2007 prescribed fire emissions, held constant at 2005 emissions level. These emission estimates are presented at the end of this section in Table H-5, along with the total emissions from the forestry and land use sector.

Table H4H5. Forestry and Land Use Flux and Reference Case Projections (MMtCO₂e)

Subsector	1990	2000	2005	2010	2020	2025
Forested Landscape (excluding soil carbon)	-3.38	-21.1	-21.1	21.049.2	-20.949.2	-20.949.2
Urban Forestry and Land Use	-14.4	-5.65	-6.23	-6.23	-6.23	-6.23
Forest Wildfires	1.35	1.15	0.16	1.00	1.00	1.00
Forest Prescribed Fires	5.70	4.14	6.66	5.70	5.70	5.70
Sector Total	-16.5	-25.6	-27.1	-20.54.4	-204.4	-204.4

Formatted Table

Subsector	1990	2000	2005	2010	2020	2025
Forested Landscape (excluding soil carbon)	0.06	-24.5	-24.5	-24.5	-24.5	-24.5
Urban Forestry and Land Use	-14.4	-5.65	-6.23	-6.23	-6.23	-6.23
Wildfires & Prescribed Burns	1.35	1.15	0.16	0.16	0.16	0.16

Formatted: Left

Sector Total	-12.1	-29.0	-30.6	-30.6	-30.6	-30.6
--------------	-------	-------	-------	-------	-------	-------

Key Uncertainties

It is important to note that there were methodological differences in the three FIA cycles (used to calculate carbon pools and flux) that can produce different estimates of forested area and carbon density. For example, the FIA program modified the definition of forest cover for the woodlands class of forestland (considered to be non-productive forests). Earlier FIA cycles defined woodlands as having a tree cover of at least 10%, while the newer sampling methods used a woodlands definition of tree cover of at least 5% (leading to more area being defined as woodland). This issue is probably of more relevance in the western US. Also, in woodland areas, the earlier FIA surveys might not have inventoried trees of certain species or with certain tree form characteristics (leading to differences in both carbon density and forested acreage). Given that the forested land in Florida is dominated by timberlands (productive forests), CCS does not believe that the definitional differences noted above have had a significant impact on the forest flux estimates provided in this report; however additional input from technical workgroup members and state foresters is needed.

Also, FIA surveys since 1999 include all dead trees on the plots, but data prior to that are variable in terms of these data. The modifications to FIA surveys are a result of an expanded focus in the FIA program, which historically was only concerned with timber resources, while more recent surveys have aimed at a more comprehensive gathering of forest biomass data. In addition, the FIA program has moved from periodic to annual inventory methods. The effect of these changes in survey methods has not been estimated by the USFS.

There was conflicting data found in two FIA databases regarding FL’s 1995 total forest area. The Carbon Calculation Tool (used to estimate forest carbon pool and flux) was based on FIADB 2.1, a version that is not up-to-date. Instead, the 1995 forest area from FIADB 3.0 was used. So the C pool and fluxes originally estimated from CCT had to be adjusted using the new 1995 forest area based on ratios. A more accurate adjustment would need to be made in the future using CCT that pulls in the most current FIA data.

Uncertainties also arise regarding the future of Florida’s forests. While the 2005-2025 projections are based on the two most recent FIA inventories (1995 and 2005), which indicate a small loss of forest area, some publications forecast greater losses in Florida’s forestlands. For example, a report published by USFS¹⁶ forecasts a loss of 8 million acres of forestland in the South between 1992 and 2020 due to urbanization, while a University of Florida report¹⁷ projects a loss of 2.7 million acres of native land in FL between 2005 and 2060. These reports point out the historic decrease in forestlands in Florida, the recent trends in population growth and

¹⁶ Wear, David N. & John G. Greis. *The Southern Resource Assessment Summary Report, Chapter 6: Land Use*. USDA Forest Service, Southern Research Station, 15 March, 2007. Retrieved July, 2008 from <http://www.srs.fs.fed.us/sustain/report/socio1/socio1.htm>.

¹⁷ Zwick, Paul D. & Margaret Carr. *Florida 2060: A Population Distribution Scenario for the State of Florida*, 15 August, 2006.

Formatted: Font: Italic

urbanization, the changing returns of the agricultural and timber industries and how they affect land use, as well as growing concerns regarding the future of Florida's forests. While these reports focus on total forest area, it's also important to note the recent increase in area of timberlands. Timberland area increased 901,000 acres in 1995 to 15.6 million acres in 2005, and increased to 16 million acres in 2006 (according to the new 2006 FIA inventory).¹⁸ These factors lead to further uncertainties in forecasting forecast carbon and flux in Florida.

Much of the urban forestry & land use emission estimates rely on national default data and could be improved with state-specific information. In particular, the carbon flux estimates associated with landfilled food and yard waste should be reviewed and revised, as data are available. Even with greater attention being paid to organics management programs in the solid waste sector (e.g. composting programs), Given the level of urbanization and population growth in FL during the 1990's, it does not seem likely that levels of landfilled food and yard waste would have fallen during this period. Additional work should be done to better integrate the results of this work in the urban landscape sector with the inventory and forecast results in the waste management sector.

¹⁸ USFS Forest Service, Forest Inventory Data Online, Florida 2006 Inventory. Retrieved July 1, 2008 from <http://www.fia.fs.fed.us/tools-data/>.